# Kenai Fjords National Park Freshwater Fish Inventory


Presented by Joe Miller Katmai National Park

Funding provided by the National Park Service Inventory and Monitoring Program


## Objectives


- Document previously undocumented freshwater fish species within KEFJ
- 1. Catostomus catostomus (longnose sucker)
- 2. Cottus aleuticus (coastrange sculpin)
- 3. Eleginus gracilis (saffron cod) \*\*\*Primarily marine
- 4. Lampetra tridentata (Pacific lamprey)
- 5. Prosopium cylindraceum (round whitefish)
- Document life history and biological characteristics of species encountered

## Study Location



•KEFJ located in Southcentral Alaska on Kenai Peninsula





#### Habitats Sampled (30+ locations)

**Addsion Lake** 

**Addison Lake Pond** 

**Beard's Hollow Lagoon** 

**Boulder Creek Lake** 

**Boulder Creek** 

**Crescent Beach Creek** 

**Crescent Beach Pond** 

Delight Lake

**Delight River** 

**Delight Estuary** 

**Desire Creek** 

**Delusion Creek** 

**Dropoff Beach Creek** 

**Exit Creek** 

**Ferrum Creek** 

James Lagoon Creek A

James Lagoon River A

**Nuka River** 

Northwestern Fjord Creek

Northwestern Fjord Pond

**Paguna Creek** 

Paguna Lagoon

**Pederson Creek** 

**Quicksand Creek** 

**Resurrection River** 

"Split Glacier Creek"

**Shelter Cove Creek** 

**Shelter Cove Lake** 

Verdant Lagoon

# Representative Habitats



# More habitat diversity







# Sampling Methods

Seines, hand nets, minnow traps, angling



# Results: low diversity, highly adaptable assemblage



Pacific staghorn sculpin





threespine stickleback



#### Common salmonids



Dolly Varden



sockeye salmon



coho salmon

#### Fishes encountered during inventory


\*\*\*All are euryhaline or anadromous\*\*\*

- 1. chum salmon
- 2. coho salmon
- 3. pink salmon
- 4. king salmon
- 5. sockeye salmon
- 6. Dolly varden
- 7. coastrange sculpin (previously undocumented)
- 8. Pacific staghorn sculpin
- 9. threespine stickleback
- 10. eulachon

#### So what??


- Fish assemblage reflects zoogeographic patterns seen elsewhere in recently deglaciated habitats:
- Colonizers are capable of using coastal dispersal routes
- 2. Fishes limited to inland dispersal are not well represented...stenohaline freshwater species not found

#### Glacial effects



During Wisconsin period,

Cordilleran ice sheet covered KEFJ



#### Conclusions

- KEFJ colonization has occurred rapidly
- Sockeye salmon, threespine sticklebacks, and coastrange sculpins found in habitats formed in past few decades.
- KEFJ is an excellent place to examine patterns of adaptation and evolution
- KEFJ freshwater fish communities are more similar to SE AK than SW AK

# Ice still covers much of the modern landscape

- Existing ice and mountainous terrain
  have limited colonization by FW fish and
  associated species diversity
- Glacial retreat is rapidly creating new aquatic habitats

# Acknowledgements:

Thanks to Aeren Heuer, Matt Grey, Glenn Dunmire, Caroline Dunmire, Tahzay Jones, Ian Martin, Joel Markis, Beth Koltun, Jess Grunblatt, Dorothy Mortenson, Ian Martin, Mike Tetreau, Shelley Hall, KEFJ staff, AK I&M staff, and everyone else! Funding Provided by NPS I&M program